

DESY 97-256

December 1997

FLAVOUR CHANGING NEUTRAL CURRENT PROCESSES AND CKM PHENOMENOLOGY

A. Ali

Deutsches Elektronen-Synchrotron DESY, Hamburg
Notkestraße 85, D-22603 Hamburg, FRG

To be published in the Proceedings of the First APCTP Workshop
Pacific Particle Physics Phenomenology, Oct. 31 - Nov. 2, 1997,
Seoul National University, Seoul, South Korea

FLAVOUR CHANGING NEUTRAL CURRENT PROCESSES AND CKM PHENOMENOLOGY

A. Ali

Deutsches Elektronen-Synchrotron DESY, Notkestrasse 85, D-22603 Hamburg, FRG

Some selected topics in flavour changing neutral current processes in B decays are discussed in the context of the Standard Model. The emphasis here is on the determination of the CKM matrix elements from the ongoing and forthcoming experiments. The topics discussed include: (i) CKM-allowed radiative decays $B \rightarrow X_s + \gamma$, (ii) CKM-suppressed radiative decays $B \rightarrow X_d + \gamma$, $B^\pm \rightarrow \rho^\pm + \gamma$ and $B^0 \rightarrow (\rho^0, \omega) + \gamma$, (iii) B^0 - $\overline{B}^0$ -mixings, and (iv) the Fleischer-Mannel bound in $B \rightarrow K\pi$ decays.

1 Determination of $|V_{ts}|$ from $\mathcal{B}(B \rightarrow X_s + \gamma)$

1.1 Experimental status

Electromagnetic penguins in B decays were first measured in 1993 by the CLEO collaboration through the exclusive decay $B \rightarrow K^* + \gamma$ ¹, followed by the measurement of the inclusive decay $B \rightarrow X_s + \gamma$ in 1994². The present CLEO measurements can be summarized as³:

$$\begin{aligned}\mathcal{B}(B \rightarrow X_s + \gamma) &= (2.32 \pm 0.57 \pm 0.35) \times 10^{-4}, \\ \mathcal{B}(B \rightarrow K^* + \gamma) &= (4.2 \pm 0.8 \pm 0.6) \times 10^{-5},\end{aligned}\tag{1}$$

which yield an exclusive-to-inclusive ratio: $R_{K^*} \equiv \Gamma(B \rightarrow K^* + \gamma)/\Gamma(B \rightarrow X_s + \gamma) = (18.1 \pm 6.8)\%$. Very recently, the inclusive radiative decay has also been reported by the ALEPH collaboration with a (preliminary) branching ratio⁴:

$$\mathcal{B}(H_b \rightarrow X_s + \gamma) = (3.29 \pm 0.71 \pm 0.68) \times 10^{-4}.\tag{2}$$

The branching ratio in (2) involves a different weighted average of the various B -mesons and Λ_b baryons produced in Z^0 decays (hence the symbol H_b) than the corresponding one given in (1), which has been measured in the decay $\Upsilon(4S) \rightarrow B^+ B^-, B^0 \overline{B}^0$.

In the context of SM, the principal interest in the inclusive branching ratio $B \rightarrow X_s + \gamma$ lies in that it determines the ratio of the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements $|V_{ts}^* V_{tb}/V_{cb}|$. Since $|V_{cb}|$ and $|V_{tb}|$ have been directly measured, the measurements (1) and (2) yield $|V_{ts}|$. The ratio R_{K^*} constrains the form factor in $B \rightarrow K^* + \gamma$. Estimates of this ratio in the QCD sum rule and lattice QCD⁵ are in agreement with the present measurements.

1.2 SM estimates of $\mathcal{B}(B \rightarrow X_s + \gamma)$ and $\mathcal{B}(H_b \rightarrow X_s + \gamma)$

The leading contribution to the decay $b \rightarrow s + \gamma$ arises at one-loop from the so-called penguin diagrams. With the help of the unitarity of the CKM matrix, the decay matrix element in the lowest order can be written as:

$$\mathcal{M}(b \rightarrow s + \gamma) = \frac{G_F}{\sqrt{2}} \frac{e}{2\pi^2} \lambda_t (F_2(x_t) - F_2(x_c)) q^\mu \epsilon^\nu \bar{s} \sigma_{\mu\nu} (m_b R + m_s L) b. \quad (3)$$

where G_F is the Fermi coupling constant, $e = \sqrt{4\pi\alpha_{\text{em}}}$, $x_i = m_i^2/m_W^2$; $i = u, c, t$ are the scaled quark mass ratios, and q_μ and ϵ_μ are, respectively, the photon four-momentum and polarization vector. Here, $F_2(x_i)$ is the Inami-Lim function derived from the (1-loop) penguin diagrams⁶ and the CKM-matrix element dependence is factorized in $\lambda_t \equiv V_{tb}V_{ts}^*$. As the inclusive decay widths of the B hadrons are proportional to $|V_{cb}|^2$, the measurement of $\mathcal{B}(B \rightarrow X_s + \gamma)$ can be readily interpreted in terms of the CKM-matrix element ratio $\lambda_t/|V_{cb}|$. The required QCD radiative and power corrections have been computed in the effective Hamiltonian approach up to next-to-leading order (NLO) logarithmic accuracy and to leading order in $1/m_b^2$ and $1/m_c^2$. The effective Hamiltonian in the SM can be written as (keeping operators up to dimension 6),

$$\mathcal{H}_{eff}(b \rightarrow s + \gamma) = -\frac{4G_F}{\sqrt{2}} \lambda_t \sum_{i=1}^8 C_i(\mu) \mathcal{O}_i(\mu), \quad (4)$$

where the operator basis, the lowest order coefficients $C_i(m_W)$ and the renormalized coefficients $C_i(\mu)$ can be seen elsewhere⁷.

Expressing the branching ratio $\mathcal{B}(B \rightarrow X_s + \gamma)$ in terms of the semileptonic decay branching ratio $\mathcal{B}(B \rightarrow X \ell \nu_\ell)$,

$$\mathcal{B}(B \rightarrow X_s \gamma) = \left[\frac{\Gamma(B \rightarrow \gamma + X_s)}{\Gamma_{SL}} \right]^{th} \mathcal{B}(B \rightarrow X \ell \nu_\ell), \quad (5)$$

the theoretical part denoted by $[...]^{th}$ has been calculated in the NLO accuracy^{8,9,10}. The power corrections have been obtained assuming that the decay $B \rightarrow X_s + \gamma$ is dominated by the magnetic moment operator O_7 . The calculated $1/m_b^2$ correction is found to be innocuous (contributing at about 1%) in $\mathcal{B}(B \rightarrow X_s + \gamma)$. The correction proportional to $1/m_c^2$, resulting from the interference of the operators O_2 and O_7 in $\mathcal{H}_{eff}(b \rightarrow s \gamma)$, has also been worked out^{11,12,13}. Expressing this symbolically as

$$\frac{\Gamma(B \rightarrow X_s + \gamma)}{\Gamma^0(B \rightarrow X_s + \gamma)} = 1 + \frac{\delta_c}{m_c^2}, \quad (6)$$

one finds¹³ $\delta_c/m_c^2 \simeq +0.03$.

Using $|V_{ts}^* V_{tb}/V_{cb}| = 0.976 \pm 0.010$, obtained from the CKM-unitarity¹⁴, and the current measurements¹⁵: $\mathcal{B}(B \rightarrow X \ell \nu_\ell) = (10.49 \pm 0.46)\%$ (at $\Upsilon(4S)$) and $\mathcal{B}(H_b \rightarrow X \ell \nu_\ell) = (11.16 \pm 0.20)\%$ (at Z^0), yields

$$\begin{aligned}\mathcal{B}(B \rightarrow X_s + \gamma) &= (3.51 \pm 0.32) \times 10^{-4}, \\ \mathcal{B}(H_b \rightarrow X_s + \gamma) &= (3.76 \pm 0.30) \times 10^{-4},\end{aligned}\quad (7)$$

to be compared with the CLEO measurement $\mathcal{B}(B \rightarrow X_s + \gamma) = (2.32 \pm 0.67) \times 10^{-4}$ and the ALEPH measurement $\mathcal{B}(H_b \rightarrow X_s + \gamma) = (3.29 \pm 0.98) \times 10^{-4}$, respectively.

Letting the CKM factor $|V_{ts}^* V_{tb}/V_{cb}|$ as a free parameter, the NLO SM-based theory and experiments have been used to determine the CKM factor, yielding⁵

$$\left| \frac{V_{ts}^* V_{tb}}{V_{cb}} \right| = 0.84 \pm 0.10. \quad (8)$$

With the CKM unitarity, one has $\left| \frac{V_{ts}^* V_{tb}}{V_{cb}} \right| \simeq |V_{cs}|$; this equality holds numerically (within present precision) if one compares eq. (8) with $|V_{cs}| = 1.01 \pm 0.18$, determined from charmed hadron decays¹⁴. Using the value of $|V_{tb}|$ measured by the CDF collaboration¹⁶, $|V_{tb}| = 0.99 \pm 0.15$, and noting that $|V_{cb}| = 0.0393 \pm 0.0028$ ¹⁷, finally yields

$$|V_{ts}| = 0.033 \pm 0.007. \quad (9)$$

This is probably as direct a determination of $|V_{ts}|$ as we will ever see, as the decay $t \rightarrow W + s$ is too daunting to measure due to the low tagging efficiency of the s -quark jet.

2 The decay $B \rightarrow X_d + \gamma$ and constraints on the CKM parameters

In close analogy with the $B \rightarrow X_s + \gamma$ case, the complete set of dimension-6 operators relevant for the processes $b \rightarrow d\gamma$ and $b \rightarrow d\gamma g$ can be written as:

$$\mathcal{H}_{eff}(b \rightarrow d) = -\frac{4G_F}{\sqrt{2}} \xi_t \sum_{j=1}^8 C_j(\mu) \hat{O}_j(\mu), \quad (10)$$

where $\xi_j = V_{jb} V_{jd}^*$ with $j = u, c, t$. The current-current operators $\hat{O}_j$, $j = 1, 2$, have implicit in them CKM factors ξ_c and ξ_u . We shall use the Wolfenstein parametrization¹⁸, in which case the matrix is determined in terms of the four parameters $A, \lambda = \sin \theta_C$, ρ and η , and one can express the above factors as :

$$\xi_u = A \lambda^3 (\rho - i\eta), \quad \xi_c = -A \lambda^3, \quad \xi_t = -\xi_u - \xi_c. \quad (11)$$

We note that all three CKM-angle-dependent quantities ξ_j are of the same order of magnitude, $O(\lambda^3)$. The branching ratio $\mathcal{B}(B \rightarrow X_d + \gamma)$ in the SM can be quite generally written as:

$$\mathcal{B}(B \rightarrow X_d + \gamma) = D_1 \lambda^2 \{ (1 - \rho)^2 + \eta^2 - (1 - \rho)D_2 - \eta D_3 + D_4 \}, \quad (12)$$

where the functions D_i depend on various parameters such as m_t, m_b, m_c, μ , and α_s . These functions were calculated in the LL approximation some time ago¹⁹ and since then their estimates have been improved²⁰, transcribing the NLO calculations done for $B \rightarrow X_s + \gamma$, thereby reducing the scale dependence in $\mathcal{B}(B \rightarrow X_d + \gamma)$ which is marked in the LL result.

To get an estimate of $\mathcal{B}(B \rightarrow X_d + \gamma)$, the CKM parameters ρ and η have to be constrained from the unitarity fits²¹. Allowing these parameters to vary over the entire allowed domain, one gets (at 95% C.L.)²⁰:

$$6.0 \times 10^{-6} \leq \mathcal{B}(B \rightarrow X_d + \gamma) \leq 3.0 \times 10^{-5}. \quad (13)$$

The present theoretical uncertainty in this rate is a factor 5, almost all of which is due to the CKM parameters, which shows that even a modest (say, $\pm 30\%$) measurement of $\mathcal{B}(B \rightarrow X_d + \gamma)$ will have a very significant impact on the CKM phenomenology. The decay $B \rightarrow X_d + \gamma$ is of interest from the point of view of direct CP violation as well. The CP asymmetry in this mode is estimated as $O(10\%)$. This should also reflect itself in the rate asymmetry in some exclusive decays such as $B^\pm \rightarrow \rho^\pm + \gamma$.

3 CKM-suppressed exclusive radiative decays $B \rightarrow V + \gamma$

Exclusive radiative B decays $B \rightarrow V + \gamma$, with $V = K^*, \rho, \omega$, are also potentially very interesting for the CKM phenomenology²². Extraction of CKM parameters from these decays would, however, involve a trustworthy estimate of the (short distance) SD- and (long distance) LD-contributions in the decay amplitudes. Present estimates based on QCD sum rules^{23,24} and soft-scattering models²⁵ yield small ($O(5\%)$) to moderate ($O(20\%)$) LD-contributions in these decays depending on the decay modes. More importantly, data on neutral and charged B mesons can be used to disentangle the LD-effects directly.

The SD-contribution in the exclusive decays $(B^\pm, B^0) \rightarrow (K^{*\pm}, K^{*0}) + \gamma$, $(B^\pm, B^0) \rightarrow (\rho^\pm, \rho^0) + \gamma$, $B^0 \rightarrow \omega + \gamma$ and the corresponding B_s decays, $B_s \rightarrow \phi + \gamma$, and $B_s \rightarrow K^{*0} + \gamma$, involve the magnetic moment operator $\mathcal{O}_7$ and the related one obtained by the change $s \rightarrow d$, $\hat{\mathcal{O}}_7$. Keeping only the SD-contribution leads to obvious relations among the exclusive decays, exemplified here by the decay rates for $(B^\pm, B^0) \rightarrow (\rho^\pm, \rho^0) + \gamma$ and

$(B^\pm, B^0) \rightarrow (K^{*\pm}, K^{*0}) + \gamma$:

$$\frac{\Gamma((B^\pm, B^0) \rightarrow (\rho^\pm, \rho^0) + \gamma)}{\Gamma((B^\pm, B^0) \rightarrow (K^{*\pm}, K^{*0}) + \gamma)} \simeq \kappa_{u,d} \left[\frac{|V_{td}|}{|V_{ts}|} \right]^2, \quad (14)$$

where $\kappa_i \equiv [F_S(B_i \rightarrow \rho\gamma)/F_S(B_i \rightarrow K^*\gamma)]^2$ involves the ratio of the transition form factors (apart from the small phase-space dependent correction) in the indicated radiative B decays; this ratio is unity in the $SU(3)$ limit. Likewise, assuming dominance of SD physics gives relations among various decay rates

$$\Gamma(B^\pm \rightarrow \rho^\pm \gamma) = 2 \Gamma(B^0 \rightarrow \rho^0 \gamma) = 2 \Gamma(B^0 \rightarrow \omega \gamma), \quad (15)$$

where the first equality holds due to the isospin invariance, and in the second $SU(3)$ symmetry has been assumed.

The LD-amplitudes in radiative B decays from the light quark intermediate states necessarily involve CKM matrix elements other than V_{td} or V_{ts} . In the CKM-suppressed decays $B \rightarrow V + \gamma$ the LD-contributions are dominantly induced by the matrix elements of the four-Fermion operators $\hat{O}_1$ and $\hat{O}_2$. Using factorization, the LD-amplitude in the decay $B^\pm \rightarrow \rho^\pm \gamma$ can be written in terms of two form factors F_1^L and F_2^L . These are found to be numerically close to each other in the QCD sum rule approach^{23,24}, $F_1^L \simeq F_2^L \equiv F_L$, hence the ratio of the LD- and the SD- contributions reduces to a number

$$\mathcal{A}_{long}/\mathcal{A}_{short} = R_{L/S}^{B^\pm \rightarrow \rho^\pm \gamma} \cdot \frac{V_{ub}V_{ud}^*}{V_{tb}V_{td}^*}. \quad (16)$$

where $R_{L/S}^{B^\pm \rightarrow \rho^\pm \gamma}$ is estimated as²⁴:

$$R_{L/S}^{B^\pm \rightarrow \rho^\pm \gamma} = -0.30 \pm 0.07. \quad (17)$$

The CKM ratio in Eq. (16) is constrained by unitarity to lie in the range²¹ $0.22 \leq V_{ub}V_{ud}^*/V_{tb}V_{td}^* \leq 0.71$, with the central value around 0.33. This would put the LD-corrections at $O(10 - 20)\%$. The analogous LD-contributions to the neutral B decays $B^0 \rightarrow \rho\gamma$ and $B^0 \rightarrow \omega\gamma$ are expected to be much smaller due to the electric charge and QCD colour factors, and are estimated to be^{24,25} $R_{L/S}^{B^0 \rightarrow \rho\gamma} \simeq R_{L/S}^{B^0 \rightarrow \omega\gamma} = 0.05$.

The relations (15), which hold on ignoring LD-contributions, get modified by including the LD-contributions. Likewise, the ratios of the CKM-suppressed and CKM-allowed decay rates given in eq. (14) get modified due to the LD contributions. The effect of the LD-contributions is estimated to be modest but not negligible in charged B decays, introducing an uncertainty of $O(15\%)$,

comparable to the uncertainty in the overall normalization due to the $SU(3)$ -breaking effects in the quantity κ_u . Neutral B -meson radiative decays are less-prone to the LD-effects, and hence one expects that to a good approximation (say, better than 10%) the ratio of the decay rates for neutral B meson obtained in the approximation of SD-dominance remains valid²²:

$$\frac{\Gamma(B^0 \rightarrow \rho\gamma, \omega\gamma)}{\Gamma(B \rightarrow K^*\gamma)} = \kappa_d \lambda^2 [(1 - \rho)^2 + \eta^2] , \quad (18)$$

where this relation holds for each of the two decay modes separately.

Finally, combining the estimates for the LD- and SD-form factors^{24,22}, and restricting the Wolfenstein parameters in the allowed range given earlier, yields: $\mathcal{B}(B^\pm \rightarrow \rho^\pm \gamma) = (1.5 \pm 1.1) \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \rho\gamma) \simeq \mathcal{B}(B^0 \rightarrow \omega\gamma) = (0.65 \pm 0.35) \times 10^{-6}$. The large range reflects to a large extent the poor knowledge of the CKM matrix elements and hence experimental measurements of these branching ratios will contribute greatly to determine the Wolfenstein parameter ρ and η . Present experimental limits (at 90% C.L.) are³: $\mathcal{B}(B^\pm \rightarrow \rho^\pm \gamma) < 1.1 \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow \rho\gamma) < 3.9 \times 10^{-5}$ and $\mathcal{B}(B^0 \rightarrow \omega\gamma) < 1.3 \times 10^{-5}$. The constraints on the parameters (ρ, η) following from them are, however, not yet competitive to the ones which follow from the CKM-unitarity and lower bound on the mass difference ratio $\Delta M_s / \Delta M_d$ in the $B_s^0 - \bar{B}_s^0$ and $B_d^0 - \bar{B}_d^0$ sectors, which we discuss next.

4 ΔM_s (and x_s), Unitarity Triangle and CP Phases

Mass differences in the $B_s^0 - \bar{B}_s^0$ and $B_d^0 - \bar{B}_d^0$ systems are dominated in the standard model by the t -quark exchange. Of the two, ΔM_d is well-measured¹⁵, $\Delta M_d = (0.464 \pm 0.018)(\text{ps})^{-1}$, which can be translated into a determination of the matrix element $|V_{td}|$, yielding²¹ $|V_{td}| = (9.0 \pm 2.6) \times 10^{-3}$, where the error reflects dominantly theoretical uncertainty on the hadronic quantity, for which $f_{B_d} \sqrt{\hat{B}_{B_d}} = (200 \pm 40) \text{ MeV}$ has been used. The mass difference ΔM_s depends quadratically on the CKM factor $|V_{ts}^* V_{tb}|$, which has been measured from the branching ratio $B \rightarrow X_s + \gamma$, yielding $|V_{ts}^* V_{tb}| = 0.033 \pm 0.007$. However, as the error on this quantity is more than a factor 2 larger than what one gets from the CKM unitarity, we use here the value obtained by unitarity, $|V_{ts}^* V_{tb}| = 0.038 \pm 0.0027$. This yields:

$$\begin{aligned} \Delta M_s &= (12.0 \pm 2.0) \frac{f_{B_s}^2 B_{B_s}}{(230 \text{ MeV})^2} (\text{ps})^{-1} , \\ x_s &= (18.2 \pm 3.0) \frac{f_{B_s}^2 B_{B_s}}{(230 \text{ MeV})^2} , \end{aligned} \quad (19)$$

where we have used $\tau(B_s) = (1.52 \pm 0.07)$ ps to calculate x_s . The choice $f_{B_s}\sqrt{\hat{B}_{B_s}} = 230$ MeV corresponds to the central value given by the lattice-QCD estimates. Allowing the coefficient to vary by $\pm 2\sigma$, and taking the central value for $f_{B_s}\sqrt{\hat{B}_{B_s}}$, gives

$$\begin{aligned} 8.0 \text{ (ps)}^{-1} &\leq \Delta M_s \leq 16.0 \text{ (ps)}^{-1} , \\ 12.2 &\leq x_s \leq 24.2 . \end{aligned} \quad (20)$$

It is difficult to ascribe a confidence level to this range due to the dependence on the unknown coupling constant factor. The present lower bound (from LEP) on ΔM_s is: $\Delta M_s > 10.0 \text{ (ps)}^{-1}$ (95% C.L.)²⁶. A useful quantity for the CKM phenomenology is the ratio

$$\frac{\Delta M_s}{\Delta M_d} = \frac{\hat{\eta}_{B_s} M_{B_s} (f_{B_s}^2 B_{B_s})}{\hat{\eta}_{B_d} M_{B_d} (f_{B_d}^2 B_{B_d})} \left| \frac{V_{ts}}{V_{td}} \right|^2 , \quad (21)$$

in which all dependence on the t -quark mass drops out, leaving the square of the ratio of CKM matrix elements, multiplied by a factor which reflects $SU(3)_{\text{flavour}}$ -breaking effects.

The present lower bound on ΔM_s from LEP and the world average of ΔM_d can be used to put a bound on the ratio $\Delta M_s/\Delta M_d$, yielding $\Delta M_s/\Delta M_d \geq 20.4$ at 95% C.L. This is significantly better than the lower bound on this quantity from the CKM fits²¹: $\Delta M_s/\Delta M_d \geq 11.4 (\xi_s/1.15)^2$, where $\xi_s \equiv (f_{B_s}\sqrt{\hat{B}_{B_s}})/(f_{B_d}\sqrt{\hat{B}_{B_d}})$ is theoretically estimated to be $\xi_s = (1.15 \pm 0.05)$. The 95% confidence limit on $\Delta M_s/\Delta M_d$ can be turned into a bound on the CKM parameter space (ρ, η) by choosing a value for the $SU(3)$ -breaking parameter ξ_s . We assume three representative values: $\xi_s^2 = 1.21, 1.32$ and 1.44 , and display the resulting constraints in Fig. 1. From this graph we see that the LEP bound now restricts the allowed ρ - η region for all three values of ξ_s^2 . We note that the bound on $\Delta M_s/\Delta M_d$ removes a large (otherwise allowed) negative- ρ region. The resulting fits of the CKM triangle taking into account this bound are necessarily asymmetric around $\rho = 0$. The best fit solutions are around $\rho = 0.11, \eta = 0.33$, as also noted by Paganini et al.²⁷.

The CP-violating asymmetries in B decays can be expressed straightforwardly in terms of the CKM parameters ρ and η . The 95% C.L. constraints on ρ and η just discussed can be used to predict the correlated ranges of the angles of the CKM unitarity triangle α, β and γ in the standard model. These correlations were worked out earlier without the $\Delta M_s/\Delta M_d$ bound^{21,7} and are being updated here. It is found that the effect of the LEP bound on $\Delta M_s/\Delta M_d$




Figure 1: Constraints in ρ - η space from the LEP bound $\Delta M_s/\Delta M_d > 20.4$. The bounds are presented for 3 choices of the SU(3)-breaking parameter: $\xi_s^2 = 1.21$ (dotted line), 1.32 (dashed line) and 1.44 (solid line). In all cases, the region to the left of the curve is ruled out. (Updated from ref. 21.)

is still marginal on the $\sin 2\beta$ - $\sin 2\alpha$ correlation; a small region in $\sin 2\beta$ below $\sin 2\beta = 0.3$ is now removed, with the (95% C.L.) range for $\sin 2\alpha$ remaining practically unconstrained with or without the LEP bound. The correlation in the angles α - γ is more susceptible to the aforementioned bound. The allowed range for this correlation also depends more sensitively on the value of the quantity B_K than is the case for the $\sin 2\beta$ - $\sin 2\alpha$ correlation. The previously allowed range^{21,7} (at 95% C.L.): $\gamma = (90 \pm 50)^\circ$ was based on using the bag parameter to have a value in the range $B_K = 0.9 \pm 0.1$. Taking a somewhat larger error on the advice of our lattice colleagues, $B_K = 0.9 \pm 0.15$, this range becomes larger. One now gets $\gamma = (90 \pm 58)^\circ$ without the LEP bound. The corresponding (95 % C.L.) range with the LEP bound is: $32^\circ \leq \gamma \leq 120^\circ$. Thus, values of the angle γ in excess of 120° are now excluded by the LEP bound on $\Delta M_s/\Delta M_d$. Moreover, the allowed range in γ is no longer symmetric around $\gamma = 90^\circ$. This is illustrated in Fig. 2, which shows the region in α - γ space allowed by the present data, for $f_{B_d}\sqrt{\hat{B}_{B_d}} = 200 \pm 40 \text{ MeV}$ and $B_K = 0.9 \pm 0.15$.

The ranges for the CP-violating rate asymmetries parametrized by $\sin 2\alpha$, $\sin 2\beta$ and $\sin^2 \gamma$ are determined at 95% C.L. to be:

$$-1.0 \leq \sin 2\alpha \leq 1.0 ,$$




Figure 2: Allowed values (in degrees) of the angle α and γ at 95% C.L. resulting from the CKM fits taking into account the LEP bound $\Delta M_s/\Delta M_d > 20.4$, with the indicated ranges of $f_{B_d}\sqrt{\hat{B}_{B_d}}$ and B_K . (Updated from ref. 21.)

$$\begin{aligned} 0.30 &\leq \sin 2\beta \leq 0.88 , \\ 0.27 &\leq \sin^2 \gamma \leq 1.0 . \end{aligned} \quad (22)$$

The CP asymmetries just discussed will be measured through rate asymmetries in non-leptonic B decays, such as $\overline{B}_d \rightarrow J/\psi K_S$ ($\sin 2\beta$), $\overline{B}_d \rightarrow \pi^+\pi^-$ ($\sin 2\alpha$) and $\overline{B}_s \rightarrow D_s^\pm K^\mp$ (or $B^\pm \rightarrow \overline{D}^\pm K^\pm$) ($\sin^2 \gamma$).

5 The Fleischer-Mannel bound in $B \rightarrow K\pi$ decays

Recently, the role of the decays $B^0 \rightarrow \pi^\mp K^\pm$ and $B^\pm \rightarrow \pi^\pm K^0$ in determining the angle γ has been emphasized by Fleischer and Mannel²⁸. Assuming that the so-called Tree (current-current) and QCD penguins are the dominant contributions in these decays, one can write the amplitudes as

$$\begin{aligned} \Gamma(B^0 \rightarrow \pi^\mp K^\pm) &\propto |A_P|^2 (1 - 2r \cos \gamma \cos \delta + r^2) , \\ \Gamma(B^\pm \rightarrow \pi^\pm K^0) &\propto |A_P|^2 , \end{aligned}$$

where $r = |A_T|/|A_P|$ is the ratio of the Tree and Penguin amplitudes, and δ is the strong phase difference involving these amplitudes. This leads to the Fleischer-Mannel relation (an average over CP-conjugate modes is implied),

$$R \equiv \frac{\Gamma(B^0 \rightarrow \pi^\mp K^\pm)}{\Gamma(B^\pm \rightarrow \pi^\pm K^0)} = 1 - 2r \cos \gamma \cos \delta + r^2 \geq \sin^2 \gamma . \quad (23)$$

From this, constraints on γ of the form

$$0^\circ \leq \gamma \leq \gamma_0 \quad \vee \quad 180^\circ - \gamma_0 \leq \gamma \leq 180^\circ \quad (24)$$

follow, where γ_0 is the maximum value of γ , which are complementary to the ones from the CKM unitarity fits discussed above. For $R < 1$, a useful bound on γ emerges, $\gamma_o^{\max} = \cos^{-1}(\sqrt{1-R})$. At present R has the value²⁹: $R = 0.65 \pm 0.40$. For the central value of R , for example, this yields $\gamma_o^{\max} = 54^\circ$, implying $0^\circ \leq \gamma \leq 54^\circ \vee 126^\circ \leq \gamma \leq 180^\circ$. One then finds that there is no overlap in the range $\gamma > 90^\circ$ (corresponding to $\rho < 0$ in Fig. 1) with the allowed range from the CKM fits, but there is significant overlap in the range $\gamma < 90^\circ$ (or $\rho > 0$ in Fig. 1), yielding $32^\circ \leq \gamma \leq 54^\circ$ for the assumed value of R . However, as R increases, the overlap with the parameter space allowed by the CKM-unitarity increases. Hence, with the present experimental error on R , no useful bound on γ follows: For the Fleischer-Mannel bound to be useful the ratio R must be significantly less than 1. With model-dependent estimates of r and δ , such as those obtained in the context of factorization models, sharper constraints follow on the CKM parameters. For example, r is estimated to lie in the range $0.15 \leq r \leq 0.30$ in such models, one then finds³⁰ that $R = 0.65 \pm 0.40$ implies $\rho \geq 0$ (at $\pm 1\sigma$). Possible dilution of the Fleischer-Mannel bounds from the neglected contributions (such as annihilation, electroweak penguins and soft rescattering) have been anticipated³¹. Some of these contributions may introduce significant uncertainties in the extraction of γ from this method. These issues will surely be studied further in future with more data available. It is too early to say if the ratio R (as well as other related ones) will make a big impact on the CKM phenomenology, but at any rate it remains a potentially interesting method to measure the otherwise daunting angle γ .

Acknowledgments

I would like to acknowledge collaboration with Hrachia Asatryan, Christoph Greub and David London on many aspects of B physics discussed here. In particular, David London provided the two updated figures shown in this report. Thanks are due to Manuel Drees and the organizing committee of the APCTP workshop in Seoul for holding an exciting meeting. The hospitality of Prof. H.S. Song at Seoul National University is gratefully acknowledged. This work is partially supported by the German-Korean scientific exchange programme DFG-446-KOR-113/72/0.

References

1. R. Ammar et al. (CLEO Collaboration), *Phys. Rev. Lett.* **71**, 674 (1993).
2. M.S. Alam et al. (CLEO Collaboration), *Phys. Rev. Lett.* **74**, 2885 (1995).

3. T. Skwarnicki, preprint HEPHY 97-03, hep-ph/9712253.
4. P.G. Colrain and M.I. Williams (ALEPH Collaboration), contributed paper # 587, EPS Conference, Jerusalem, August 1997.
5. A. Ali, DESY 97-192, hep-ph/9709507, to be published in Proc. of the Seventh Int. Symp. on Heavy Flavors, Santa Barbara, Calif., 1997.
6. T. Inami and C.S. Lim, *Prog. Theor. Phys.* **65**, 297 (1981).
7. A. Ali, *Acta Physica Polonica B* **27**, 3529 (1996).
8. K. Chetyrkin, M. Misiak and M. Münz, *Phys. Lett. B* **400**, 206 (1997).
9. C. Greub and T. Hurth, hep-ph/9708214.
10. A.J. Buras, A. Kwiatkowski and N. Pott, hep-ph/9707482.
11. M.B. Voloshin, *Phys. Lett. B* **397**, 275 (1997).
12. A. Khodjamirian et al., *Phys. Lett. B* **402**, 167 (1997); Z. Ligeti, L. Randall and M.B. Wise, *Phys. Lett. B* **402**, 178 (1997); A.K. Grant et al., *Phys. Rev. D* **56**, 3151 (1997).
13. G. Buchalla, G. Isidori and S.J. Rey, hep-ph/9705253.
14. R.M. Barnett et al. (Particle Data Group), *Phys. Rev. D* **54**, 1 (1996).
15. M. Feindt, plenary talk, EPS Conference, Jerusalem, August 1997.
16. M. Narain, in Proc. of the Seventh Int. Symp. on Heavy Flavors, Santa Barbara, Calif., 1997.
17. L. Gibbons, in Proc. of the Seventh Int. Symp. on Heavy Flavors, Santa Barbara, Calif., 1997.
18. L. Wolfenstein, *Phys. Rev. Lett.* **51**, 1845 (1983)
19. A. Ali and C. Greub, *Phys. Lett. B* **287**, 191 (1992).
20. A. Ali, H.M. Asatrian, and C. Greub, DESY 97-255 (1997).
21. A. Ali and D. London, *Nucl. Phys. B*(Proc. Suppl.) **54A**, 297 (1997).
22. A. Ali, V.M. Braun and H. Simma, *Z. Phys. C* **63**, 437 (1994).
23. A. Khodjamirian, G. Stoll, and D. Wyler, *Phys. Lett. B* **358**, 129 (1995).
24. A. Ali and V.M. Braun, *Phys. Lett. B* **359**, 223 (1995).
25. J.F. Donoghue, E. Golowich and A.A. Petrov, *Phys. Rev. D* **55**, 2657 (1997).
26. H.G. Moser, talk at the Heavy Flavor Workshop, Rostock, 1997.
27. P. Paganini, F. Parodi, R. Roudeau and A. Stochi, hep-ph/9711261.
28. R. Fleischer and T. Mannel, preprint TTP97-17 (hep-ph/9704423).
29. R. Godang et al. (CLEO Collaboration), preprint CLNS 97-1522, CLEO 97-27 (1997).
30. A. Ali and C. Greub, hep-ph/9707251, to appear in Phys. Rev. D.
31. J.M. Gerard and J. Weyers, preprint UCL-IPT-97-18 (hep-ph/9711469); M. Neubert, preprint CERN-TH/97-342 (hep-ph/9712224); A.F. Falk et al., preprint JHU-TIPAC-97018 (hep-ph/9712225).